

The Kohn-Luttinger Effect in Gauge Theories

T. Schäfer

Department of Physics, North Carolina State University, Raleigh, NC 27695

Kohn and Luttinger showed that a many body system of fermions interacting via short range forces becomes superfluid even if the interaction is repulsive in all partial waves. In gauge theories such as QCD the interaction between fermions is long range and the assumptions of Kohn and Luttinger are not satisfied. We show that in a $U(1)$ gauge theory the Kohn-Luttinger phenomenon does not take place. In QCD attractive channels always exist, but there are cases in which the primary pairing channel leaves some fermions ungapped. As an example we consider the unpaired fermion in the 2SC phase of QCD with two flavors. We show that it acquires a very small gap via a mechanism analogous to the Kohn-Luttinger effect. The gap is too small to be phenomenologically relevant.

I. INTRODUCTION

The conventional picture of superconductivity in cold Fermi liquids is that attractive interactions between fermions lead to a pairing instability that destroys the Fermi surface. Bardeen, Cooper, and Schrieffer showed that this instability is present for any attractive interaction, no matter how weak [1]. This would seem to imply that a Fermi liquid with purely repulsive interactions is stable and has a Fermi surface.

Kohn and Luttinger [2] showed that in the case of fermions interacting via short range forces this is not correct. There is a pairing instability even if the interaction is repulsive in all partial wave channels. The resulting critical temperature is typically very small, but the Kohn-Luttinger effect may well play a role in p -wave pairing in liquid ^3He [3, 4] and cold atomic gases [5] as well as d -wave pairing in high T_c superconductivity.

In this paper we study whether the Kohn-Luttinger effect also takes place in gauge theories. This study is part of a broader program to understand to what extent results from Fermi liquid theory can be extended to dense systems of fermions interacting via long range gauge forces. Non-Fermi liquid effects in the specific heat and the fermion dispersion relation were studied in [6, 7, 8, 9, 10, 11]. The specific question whether repulsive interactions in gauge theories can lead to superfluidity is of interest in connection with effective $U(1)$ gauge theories that arise in condensed matter systems. In non-abelian gauge theories attractive channels always exist. However, there are a number of superconducting phases in which some of the fermions remain ungapped and the residual interactions between these excitations are repulsive. A simple example is color superconductivity in QCD with two flavors. In this case up and down quarks with two different colors form an s -wave superconductor whereas up and down quarks of the third color remain gapless. Gapless fermions also appear in the gapless color-flavor-locked (CFL) phase [12].

II. SHORT RANGE FORCES

Kohn and Luttinger considered short range interactions characterized by a BCS potential $V(x)$. This potential describes particle-particle scattering $\vec{p}_1 + (-\vec{p}_1) \rightarrow \vec{p}_3 + (-\vec{p}_3)$ with $|\vec{p}_1| = |\vec{p}_3| = p_F$ and $x = \hat{p}_1 \cdot \hat{p}_3$ is the scattering angle. The partial wave amplitudes are

$$V_l = \int dx P_l(x) V(x), \quad (1)$$

where $P_l(x)$ is the Legendre polynomials of order l . If one of the partial wave amplitudes is attractive then s -channel rescattering between the particles will lead to a logarithmic growth of the interaction as the energy of the pair goes to zero. The system becomes superfluid when the effective interaction reaches a Landau pole. If all V_l are repulsive then the effective interaction goes to zero logarithmically near the Fermi surface.

Kohn and Luttinger observed that in this case t -channel particle-hole exchanges cannot be neglected. They note that if the BCS amplitude and all its derivatives are analytic then the tree level partial wave amplitudes have to be exponentially small, $V_l \sim e^{-l}$, as $l \rightarrow \infty$. This implies that perturbative corrections to the effective interaction can be important even if the coupling is small. In particular, if perturbative corrections are non-analytic in the scattering angle x then they will dominate over the tree level amplitude for sufficiently large l . The t -channel particle-hole bubble shown in Fig. 1 is

$$\Pi(\nu=0, q) = \frac{mk_F}{4\pi^2} \left\{ -1 + \frac{1}{q} \left(1 - \frac{q^2}{4} \right) \log \left(\frac{2-q}{2+q} \right) \right\}, \quad (2)$$

where $\nu = k_0$ is the energy transfer and $q = |\vec{k}|/k_F$ is the momentum transfer in units of k_F . This expression has a logarithmic singularity at $q = 2$ which corresponds to $x = -1$. The contribution of the singularity to the BCS amplitude is

$$\delta V(x) = -[V(-1)]^2 \frac{mk_F}{16\pi^2} (1+x) \log(1+x). \quad (3)$$

The correction to the partial wave amplitudes can be




FIG. 1: The upper panel shows the direct interaction and the t -channel particle-hole correction in a theory with short range interactions. The lower panel shows the direct interaction in a gauge theory. In this case, t -channel particle-hole bubbles have to be resummed at leading order.

determined by using the generating functional

$$P_l(x) = \frac{1}{2^l l!} \frac{d^l}{dx^l} \left[(x^2 - 1)^l \right] \quad (4)$$

and integrating by parts l times. We find

$$\delta V_l = (-1)^{l+1} \frac{mk_F}{4\pi^2} \frac{[V(-1)]^2}{l^4} \quad (5)$$

which is attractive for even l and falls off as a power of l . This implies that even if all V_l are repulsive there is a critical l_{crit} such that $V_l + \delta V_l$ is attractive for l even and $l > l_{crit}$.

In nuclear physics the screening correction is known as the “induced interaction”. Consider a short range potential with a strength adjusted to give the the scattering length a . If a is negative then the s -wave interaction is attractive and in BCS theory we find a gap

$$\Delta = \frac{8E_F}{e^2} \exp\left(-\frac{\pi}{2k_F|a|}\right), \quad (6)$$

where k_F is the Fermi momentum and E_F is the Fermi energy. The screening correction reduces the magnitude of the gap by a factor $(4e)^{1/2} \simeq 2.2$ [13]. If the a is positive then the interaction is repulsive and no s -wave pairing takes place. However, the interaction is attractive in higher partial waves. The d -wave gap is of order

$$\Delta \sim E_F \exp\left(-\frac{\pi^2}{(2k_F a)^2} \frac{105}{2(8 - \log(2048))}\right) \quad (7)$$

Higher order corrections to this result were studied in [3, 4].

III. SUPERCONDUCTIVITY IN GAUGE THEORIES

In gauge theories the interaction is long range and the particle-hole screening correction has to be included at leading order even if the angular momentum l is small.

The screened electric and magnetic gauge boson propagators are given by [14, 15]

$$iD_{00}(\omega, \vec{k}) = \frac{1}{\vec{k}^2 + 2m^2}, \quad (8)$$

$$iD_{ij}(\omega, \vec{k}) = \frac{\delta_{ij} - \hat{k}_i \hat{k}_j}{\vec{k}^2 - i\frac{\pi}{2}m^2\omega/|\vec{k}|}, \quad (9)$$

where m^2 is the dynamically generated gauge boson mass and we have assumed that $|\omega| \ll |\vec{k}|$. In perturbation theory $m^2 = g^2 N_f \mu^2 / (4\pi^2)$ where g is the gauge coupling, μ is the baryon chemical potential and N_f is the number of fermions in the fundamental representation. The electric interaction is completely screened. At low energy multipole moments of the interaction are independent of energy and decrease exponentially with l . The magnetic interaction, on the other hand, is only dynamically screened. Multipole moments diverge as a logarithm of energy and the coefficient of $\log(\omega)$ is independent of l . We find [16]

$$f_l(\omega) = -c_A \frac{2g^2}{3} \log(\omega), \quad (10)$$

where c_A is a symmetry factor that depends on the channel and the gauge group. For color anti-symmetric quark pairing in QCD $c_A = -(N_c + 1)/(2N_c)$. The renormalization group flow of the interaction equ. (10) was first studied by Son [16]. The value of the s -wave gap is

$$\Delta_0 \simeq 2\Lambda_{BCS} \exp\left(-\frac{\pi^2 + 4}{8}\right) \exp\left(-\frac{3\pi^2}{\sqrt{2}g}\right). \quad (11)$$

where the scale is given by $\Lambda_{BCS} = 256\pi^4 (2/N_f)^{5/2} g^{-5} \mu$ and we have set $N_c = 3$. The coefficient of $1/g$ in the exponent is determined by equ. (10) and is independent of l . The pre-exponential factor is sensitive to electric gauge boson exchanges and fermion self-energy corrections and depends on l . The dependence on l can be determined by expanding the magnetic and electric gauge boson exchange amplitudes around the forward direction. The result is $\Delta_l = \exp(3c_l) \Delta_0$ with [17, 18]

$$c_l = \int \frac{dx}{1-x} (P_l(x) - 1). \quad (12)$$

The coefficients c_l can be computed using the recursion relation for the Legendre polynomials. We find

$$c_l = -2 \sum_{n=1}^l \frac{1}{n} \xrightarrow{l \rightarrow \infty} -2 \log(l) \quad (13)$$

which implies that the gap scales as $\Delta_l \sim \Delta_0/l^6$ and becomes very small for $l \gg 1$. We should note, however, that the result is reliable only if $\log(l) < 1/g$. When this inequality is violated we cannot expand the scattering amplitude around $x = 1$. In practice, this is already the case for $l \sim 2$.




FIG. 2: Scaled gauge boson exchange multipole amplitudes $\tilde{f}_l(\tilde{\omega})$ as a function of the dimensionless energy variable $\tilde{\omega}$. The figure shows the $l = 5, 10, 20, 40$ multipoles. For QCD at $\mu = 500$ MeV the energy scale corresponds to roughly 1 GeV.

This suggests that the behavior of the scattering amplitude changes when $\log(l) \sim 1/g$, and that something analogous to the Kohn-Luttinger effect may take place. In order to study this question we compute the partial wave amplitudes without relying on an expansion around $x = 1$. The transverse gauge boson exchange contribution is given by

$$f_l(\omega) = c_R g^2 \tilde{f}_l(\tilde{\omega}), \quad \tilde{\omega} = \frac{\pi}{4\sqrt{2}} \frac{m^2}{\mu^2} \left(\frac{\omega}{\mu} \right) \quad (14)$$

where c_R is a color factor and

$$\tilde{f}_l(\tilde{\omega}) = \int dx \frac{P_l(x)}{1 - x + \tilde{\omega}/\sqrt{1-x}}. \quad (15)$$

In Fig. 2 we show the function $\tilde{f}_l(\tilde{\omega})$ for several different values of l . We observe that for every value of $\tilde{\omega}$ the scattering amplitude changes sign if l is larger than some critical value. Also, for every value of l ($l \geq 2$) the amplitude changes sign for $\tilde{\omega} > \tilde{\omega}_0(l)$. The function $\tilde{f}_l(\tilde{\omega})$ exhibits a very simple scaling behavior $\tilde{f}_l(\tilde{\omega}) = F(\tilde{\omega}_0(l)/\tilde{\omega})$ with $\tilde{\omega}_0(l) \sim l^{-\alpha}$ and $\alpha \simeq 2/\log(2)$.

The figure shows that even if the interaction in the forward direction is repulsive higher multipole amplitudes are attractive for almost all energies. The question is whether this is sufficient in order to produce superconductivity. In the appendix we discuss conditions under which interactions that are repulsive in some energy regime induce pairing instabilities. We show that if the interaction is repulsive below some energy ω_0 then a necessary condition for pairing is that the gap Δ is larger than ω_0 . An upper bound on the gap can be obtained by assuming that $\Delta(\omega)$ is independent of ω . In this case we find

$$1 = -\frac{g^2 c_R}{8\pi^2} \int_{\Delta}^{\infty} \frac{d\omega}{\omega} \tilde{f}_l(\omega) \quad (16)$$

and consistency requires that $\Delta > \omega_0(l)$. This means that the integral on the RHS can be bounded by replacing the lower limit by $\omega_0(l)$. This integral is only weakly dependent on l and quickly approaches a finite limiting value, see Fig. 2. Numerically we find

$$\lim_{l \rightarrow \infty} \int_{\omega_0(l)}^{\infty} \frac{d\omega}{\omega} \tilde{f}_l(\omega) = -0.107. \quad (17)$$

This implies that in the limit of large l a gap can only appear if $0.107 g^2 c_R / (8\pi^2) > 1$. We can ask if this result is modified by higher order perturbative corrections. If $\omega_0(l)$ is smaller than the scale of non-Fermi liquid effects in the normal phase, $\omega_{nfl} = m \exp(-9\pi^2/g^2)$, then self energy corrections to the fermion propagator have to be taken into account. These corrections further reduce equ. (17). Higher order contributions to the vertex scale as $\omega^{1/3}$ and can be neglected [11]. We conclude that in the weak coupling limit superconductivity does not take place in repulsive channels. Indeed, judging from the weak coupling result, superconductivity is not likely even for large values of the coupling.

IV. CONCLUSIONS

Our results are directly applicable to a $U(1)$ gauge theory at finite density. In that case there are no attractive channels and our results imply that superconductivity does not take place. Of course, if the density is very small or the coupling is very large then the system may exhibit other instabilities, for example towards ferromagnetism or crystalline order [19].

In QCD attractive channels always exist. However, there are cases in which the primary pairing channel leaves some modes unpaired. The simplest example is the 2SC phase of $N_f = 2$ QCD. In this case up and down quarks with two different colors are paired but quarks of the third color are unpaired. At energies below the primary pairing gap Δ these quarks interact via the exchange of the hypercharge gluon of the $SU(3)$ gauge group. This gluon is part of the broken sector of the gauge group, and as a consequence the magnetic interaction is screened. The momentum dependent screening mass was computed in [20, 21]. The result is

$$\text{Re } \Pi_T(k_0, k) = \frac{\pi^2}{4} m^2 \frac{\Delta}{k} \quad (k_0 \ll \Delta, k \gg \Delta). \quad (18)$$

For $k < \Delta$ the screening mass goes to a constant. We observe that this result has the same structure as the Landau damping term except that the self energy is real rather than imaginary and the energy is replaced by the gap. As a consequence the partial wave amplitudes are identical to the results shown in Fig. 2 for $\omega > \Delta$ and become constant for $\omega < \Delta$. This means that pairing takes place in the partial wave for which $\omega_0 \sim \Delta$, so $l \sim (m/\Delta)^{1/\alpha}$. This is parametrically large, but for values of the gap that are of physical interest l could be as small as $l = 2$.

Finally we can try to estimate the magnitude of the pairing gap. The scale for the secondary gap Δ_R is set by the primary gap Δ . The multipole amplitude $f_l(\omega)$ is independent of energy for $\omega < \Delta$ and the dependence of the gap on the coupling constant is given by the BCS result, $\Delta_R \sim \Delta \exp(-c/g^2)$. The constant $c = 8\pi^2/(c_R|f_l(\Delta)|)$ is determined by the color factor c_R and the partial wave amplitude at the scale set by the primary gap. Using $c_R = 1/3$ for the ungapped quark in the 2SC phase and $|f_l(\Delta)| \leq 0.07$ from Fig. 2 we get $c \simeq 3300$. This means that the gap is completely negligible, even if the coupling is strong. In practice the most likely mechanism for generating a secondary gap is the attractive instanton interaction in the spin-one channel $\langle uC\bar{a}d \rangle$ discussed in [22]. This mechanism requires strong coupling, since perturbative interactions in this channel are repulsive.

Acknowledgments: I would like to thank I. Shovkovy for a useful discussion. This work was supported by US Department of Energy grants DE-FG02-03ER41260.

APPENDIX A: ENERGY DEPENDENT INTERACTIONS

In this appendix we study how superconductivity can arise in microscopic models that contain an effective interaction which depends on energy and is repulsive near the Fermi surface. Consider a gap equation of the type

$$\Delta(\omega) = - \int d\nu \frac{V(\omega, \nu) \Delta(\nu)}{\sqrt{\nu^2 + \Delta(\nu)^2}}. \quad (\text{A1})$$

Clearly, this equation has a non-trivial solution if the potential is attractive at low energy. However, non-trivial solutions can also arise if the potential is repulsive. Consider a potential that is repulsive everywhere, but less repulsive for $\omega < \omega_D$ [23],

$$V(\omega, \nu) = \begin{cases} V_R - V_A & |\omega - \nu| < \omega_D, \\ V_R & |\omega - \nu| > \omega_D. \end{cases} \quad (\text{A2})$$

A natural ansatz for the gap is

$$\Delta(\omega) = \begin{cases} \Delta_1 & |\omega| < \omega_D \\ \Delta_2 & |\omega| > \omega_D \end{cases} \quad (\text{A3})$$

which leads to two coupled equations

$$\Delta_1 = -(V_R - V_A) \int_0^{\omega_D} d\omega \frac{\Delta_1}{\sqrt{\omega^2 + \Delta_1^2}} - V_R \int_{\omega_D}^{\Lambda} d\omega \frac{\Delta_2}{\sqrt{\omega^2 + \Delta_2^2}} \quad (\text{A4})$$

$$\Delta_2 = -V_R \int_0^{\omega_D} d\omega \frac{\Delta_1}{\sqrt{\omega^2 + \Delta_1^2}} - V_R \int_{\omega_D}^{\Lambda} d\omega \frac{\Delta_2}{\sqrt{\omega^2 + \Delta_2^2}} \quad (\text{A5})$$

This system of equations has a non-trivial solution if the effective interaction V_{eff}

$$V_{eff} = V_A - \frac{V_R}{1 + V_R \log(\Lambda/\omega_D)} \quad (\text{A6})$$

is positive. This result has a simple interpretation from the point of view of the renormalization group. The repulsive interaction is screened as the energy is lowered from the cutoff scale Λ to the low energy scale ω_D . If V_A is larger than the screened V_R then V_{eff} is attractive. The gaps are $\Delta_1 = 2\omega_D \exp(-1/V_{eff})$ and

$$\Delta_2 = -\Delta_1 \frac{V_R}{V_{eff}(1 + V_R \log(\Lambda/\omega_D))}. \quad (\text{A7})$$

The other situation of interest is the case of a potential that is repulsive near the Fermi surface, but attractive at higher energies

$$V(\omega, \nu) = \begin{cases} V_R & |\omega - \nu| < \omega_D, \\ -V_A & |\omega - \nu| > \omega_D. \end{cases} \quad (\text{A8})$$

In this case the scale is set by the gap Δ for the purely attractive case $V_R = -V_A$. If $\Delta < \omega_D$ then there is no non-trivial solution for $V_R \geq 0$. If, on the other hand, $\Delta > \omega_D$ then there is a non-vanishing gap even if $V_R > 0$ provided that V_R remains perturbative, $V_R < 1$. Again this result is easy to understand in the context of the renormalization group. If the RG reaches a Landau pole above $\omega = \omega_D$ then the repulsive low energy potential does not modify the evolution.

-
- [1] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Phys. Rev. 108, 1175 (1957).
 - [2] W. Kohn and J. M. Luttinger, Phys. Rev. Lett. 15, 524 (1965).
 - [3] M. A. Baranov, A. V. Chubukov, M. Y. Kagan, Int. J. of Mod. Phys. B 6, 2471 (1992).
 - [4] D. V. Efremov, M. S. Mar'enko, M. A. Baranov, M. Yu. Kagan, Sov. Phys. JETP 90 861 (2000).
 - [5] A. Bulgac, M. M. Forbes and A. Schwenk, Phys. Rev. Lett. 97, 020402 (2006).
 - [6] T. Holstein, A. E. Norton, P. Pincus; Phys. Rev. B8, 2649 (1973).
 - [7] G. Baym and S. A. Chin, Nucl. Phys. A 262, 527 (1976).
 - [8] J. Polchinski, Nucl. Phys. B 422, 617 (1994).
 - [9] S. Chakravarty, R. E. Norton and O. F. Syljuasen, Phys. Rev. Lett. 75, 1423 (1995).
 - [10] A. Ipp, A. Gerhold and A. Rebhan, Phys. Rev. D 69, 011901 (2004); A. Gerhold and A. Rebhan, Phys. Rev. D 71, 085010 (2005).
 - [11] T. Schäfer and K. Schwenzer, Phys. Rev. D 70, 054007 (2004); preprint, hep-ph/0512309.
 - [12] M. Alford, C. Kouvaris and K. Rajagopal, Phys. Rev.

- Lett. **92**, 222001 (2004).
- [13] L. P. Gorkov and T. K. Melik-Barkhudarov, Sov. Phys. JETP **13**, 1018 (1961).
 - [14] J. Kapusta, Finite temperature field theory, Cambridge University Press, Cambridge (1989).
 - [15] M. LeBellac, Thermal Field Theory, Cambridge University Press, Cambridge (1996).
 - [16] D. T. Son, Phys. Rev. D **59**, 094019 (1999).
 - [17] T. Schäfer, Phys. Rev. D **62**, 094007 (2000).
 - [18] W. E. Brown, J. T. Liu and H. c. Ren, Phys. Rev. D **62**, 054016 (2000).
 - [19] F. Bloch, Z. Phys. 57, 549 (1929); E. P. Wigner, Phys. Rev. 46, 1002 (1934); D. M. Ceperley, B. J. Alder Phys. Rev. Lett. 45, 566 (1980).
 - [20] D. H. Rischke, Phys. Rev. D **62**, 034007 (2000).
 - [21] D. H. Rischke and I. A. Shovkovy, Phys. Rev. D **66**, 054019 (2002).
 - [22] M. G. Alford, K. Rajagopal and F. Wilczek, Phys. Lett. B **422**, 247 (1998).
 - [23] C. Nayak, Lecture Notes, unpublished.